

A High-Resolution Study of η Carinae's Outer Ejecta

Kerstin Weis¹

*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121
 Bonn, Germany and University of Minnesota, 116 Church Street SE,
 Minneapolis, MN 55455, USA*

Michael F. Corcoran

*Laboratory for High Energy Astrophysics, Goddard Space Flight Center,
 Greenbelt, MD 20771, USA*

Kris Davidson & Roberta M. Humphreys

*University of Minnesota, 116 Church Street SE, Minneapolis, MN
 55455, USA*

Abstract. η Carinae is a very luminous and unstable evolved star. Outflowing material ejected during the star's giant eruption in 1843 surrounds it as a nebula which consists of an inner bipolar region (the *Homunculus*) and the *Outer Ejecta*. The outer ejecta is very filamentary and shaped irregularly. Kinematic analysis, however, shows a regular bi-directional expansion despite of the complex morphology. Radial velocities in the outer ejecta reach up to 2000 km s^{-1} and give rise to X-ray emission first detected by ROSAT. We will present a detailed study of the outer ejecta based on HST images, high-resolution echelle spectra for kinematic studies, images from CHANDRA/ACIS and HST-STIS spectra.

1. Morphology, kinematics and the soft X-ray emission

The outer part of the nebula around η Carinae contains a countless number of knots, bullets and filaments and manifests the so-called outer ejecta (diameter $60''$ or 0.67 pc). A kinematic analysis using high-resolution echelle spectra showed that the outer ejecta expand with velocities between -600 km s^{-1} and $+600 \text{ km s}^{-1}$ on average (e.g. Meaburn et al. 1996, Weis 2001a,b). Various filaments nevertheless show much higher radial velocities—e.g. $\sim 2000 \text{ km s}^{-1}$ (Weis 2001a,b). It is still unclear what triggered the outburst in 1843 and which mechanism not only formed η Carinae's nebula but also the amazingly high expansion velocities. With velocities that high, X-ray emission from η Carinae is expected and was indeed detected (e.g. Cheblowski et al. 1984, Corcoran et al. 1997). With ROSAT and CHANDRA we are now able to separate between a harder,

¹Feodor-Lynen-fellow, Alexander-von-Humboldt foundation




Figure 1. In grayscale the optical (F658N) HST image (WF chip) is shown and in contours the emission in the 0.6–1.2 keV band as detected with the CHANDRA ACIS S3. Numbers indicate the expansion velocities of certain areas, underlined numbers are negative velocities, for better illustration the font size increases for higher velocities.

nearly point like emission from the central source and an extended softer emission from η Carinae's outer ejecta which is roughly hook shaped (e.g. Weis et al. 2001). Interestingly, an overlay of the optical image and the X-ray emission shows only a few correlations between the optically emitting gas and the hot X-ray gas, see Fig. 1. A much stronger conformance was found in comparison with the radial expansion velocities. The expansion velocities are derived from our optical echelle spectra ($\text{FWHM} \sim 14 \text{ km s}^{-1}$) and overplotted in Fig. 1 with negative velocities underlined. Areas with higher X-ray emission show in general higher expansion velocities. The faster the gas is expanding the more intense is the X-ray emission. The extended soft X-ray emission from η Carinae and especially the morphology of the X-ray nebula can therefore be explained by faster moving filaments which form X-ray emitting shocks. The temperature of the gas (using thermal equilibrium models) is $\sim 0.65 \text{ keV}$ (Weis et al. 2002, in prep.) indicating post-shock velocities of around 750 km s^{-1} in very good agreement with the detected expansion velocities of the bulk of the gas.

References

- Chlebowski, T., Seward, F. D., Swank, J., Szymkowiak, A., 1984, *ApJ*, 281, 665
 Corcoran, M. F., Ishibashi, K., Swank, J. H., et al. 1997, *Nature*, 390, 587
 Meaburn, J., Boumis, P., Walsh, J. R., et al. 1996, *MNRAS*, 282, 1313
 Weis, K., 2001a, in η Carinae and other mysterious stars, ed. T. Gull, S. Johanson, & K. Davidson, ASP Conf. 242, 129
 Weis, K., 2001b, in *Reviews in Modern Astronomy* 14, Springer-Verlag, ed.: Schielicke, R.E, ISBN 3-9805176-4-0, 261
 Weis, K., Duschl, W.J., Bomans, D.J., 2001, *A&A*, 367, 566